

SPATIOTEMPORAL CHAOS DUE TO SPIRAL WAVES CORE EXPANSION AND ITS ELIMINATION IN EXCITABLE MEDIA

H. Sabbagh

American University of Iraq
AUI-S
Iraq
leosabbagh@excite.com

Abstract

In the framework of the Fitzhugh Nagumo kinetics and the oscillatory recovery in excitable media, we present a new type of meandering of the spiral waves, which leads to spiral break up and spatiotemporal chaos. The tip of the spiral follows an outward spiral-like trajectory and the spiral core expands in time. This type of destabilization of simple rotation is attributed to the effects of curvature and the wave-fronts interactions in the case of oscillatory damped recovery to the rest state. This model offers a new route to and caricature for cardiac fibrillation, and when we apply the feedback resonant drift method, for defibrillation all wave activity gets eliminated at the unexcitable boundaries.

Key words

Spiral break up, spatiotemporal chaos, resonant drift.

1 Introduction

Rotating spiral waves are ubiquitous in excitable media. They have been observed in chemical reactive solutions [Kapral and Showalter, 1994], [Imbuhl and Ertl, 1995], in slime-mold aggregates [Murray, 1989] and most importantly in cardiac muscle [Chaos, Special issue on dynamics in cardiac tissue, 2002].

Such wave patterns have been studied using reaction-diffusion equations models. For some values of the system control parameters, they undergo simple rigid rotation around a circular core. However, as the control parameter is varied, the spiral tip deviates from circular trajectories [Winfree, 1987], [Meron, 1991b], [Zykov, 1986], [Zykov, 1987], [Meron, 1989], [Meron, 1991a], [Tyson and Keener, 1988]. This non-steady rotation is known as meandering and it has been observed essentially in chemical systems such as in the Belousov-Zhabotinsky (BZ) reaction [Zykov, Steinbock and Muller, 1994]. Experiments with this reaction have also demonstrated spiral breakup [Nagy-Ungvari

and Muller, 1994], [Markus and Stavridis, 1994]. This later is of interest in cardiology since it is the prelude to cardiac fibrillation, the commonest cause of sudden cardiac death [Panfilov and Hogeweg, 1995], [Winfree, 1994], and has been observed in models that show wave trains spatiotemporal instabilities [Karma, 1994], [Courtemanche, Glass, and Kenner, 1993]. It is characterized by spatiotemporally chaotic or irregular wave patterns in excitable media and remains a challenging problem in nonlinear science.

We present in this paper a new type of meandering leading to spiral breakup and offering a new route to spatiotemporal irregularity or chaos in excitable media. Spiral core expansion occurs here as the spiral free end or tip follows an outward motion along a path that looks itself like a spiral. We have previously reported some earliest results about this type of meandering [Sabbagh, 2009], and in this paper we account for this type of meandering as due to core expansion and we identify it as a new route to spiral break up and then spatiotemporal irregularity. In addition, we apply the feedback resonant drift method to eliminate the wave activity as a caricature of eliminating defibrillation. This core expansion was previously expected by the theory of non-local effects [Meron, 1991b], [Meron, 1989], [Meron, 1991a], and was attributed to effect of curvature on the velocity of propagation coupled to the effects of the interaction of successive wave-fronts due to refractoriness. The dependence of the normal velocity of propagation on curvature is given by $v = v_0 - kD$, where k is the local curvature, D is the diffusion coefficient and v_0 is the plane wave velocity of propagation [Tyson and Keener, 1988]. Due to this velocity gradient, small wavelength perturbations on the segments away from the tip would decay, which would stabilize wave propagation away from the tip and maintains the rotational motion of the spiral. On the contrary, perturbations straightening a small segment containing the tip would reduce curvature, and consequently the normal veloc-

ity of wave propagation is enhanced as the gradient of the normal velocity becomes weaker. This means that the tip would have a less tendency to curl but it tends to advance further. Therefore, further straightening of this segment containing the tip is expected. Thus, the spiral tip undergoes an outward forward motion instead of simple rigid rotation. If the recovery is non-oscillatory but monotonic, this destabilizing effect of curvature would be counteracted by the repulsive wave-front interaction due to the refractory period imposed on the medium after the passage of the preceding wave. In that case, circular rigid rotation would be sustained.

This outward motion of the tip along a spiraling trajectory was predicted by Ehud Meron in his theory of non-local effects [Meron, 1991b], [Meron, 1991a]. He proposed an approximate spiral wave solution of the reaction diffusion system in the form of a superposition of solitary wave-fronts parallel to each other, and then derived an evolution equation using a singular perturbation approach. The numerical solution of this equation, for the case of an oscillatory recovering excitable medium, was a spiral wave whose core expands in time and whose tip moves itself along a spiraling path. However, no observation of this type of spiral wave meandering and core expansion was obtained by Meron in reaction diffusion systems.

2 The Model

Here, we present a new model showing for the first time this predicted core expansion. We use a modified Barkley’s model [Barkley, 1991], [Barkley, Kness and Tuckerman, 1990] given by:

$$\begin{aligned} \partial u / \partial t &= \frac{1}{\epsilon} u(1-u)[u - ((b+v)/a)] + \nabla^2 u, \\ \partial v / \partial t &= u^3 - v, \end{aligned} \quad (1)$$

where u and v are the excitation and recovery variables respectively. The parameter b determines the excitation threshold. The inverse of ϵ , characterizing the abruptness of excitation, determines the recovery time. In the standard Barkley’s model where the local kinetics in the second equation is given by $(u - v)$, propagation cannot be maintained upon increasing ϵ . Here propagation is maintained due to the delay in the production of v .

Numerical simulations were performed on square grids using the explicit Euler integration method with a 9-point neighborhood of the Laplacian and no-flux boundary conditions. The space and time steps are respectively $dx = 0.51$ and $dt = 0.052$. Fig. 1 illustrates the nullclines. The time signals of the two variables when the threshold of excitation is exceeded are shown in Fig. 2. In the standard Barkley’s model where the local kinetics in the second equation of (1) is given by $(u - v)$, propagation cannot be maintained upon increasing ϵ . Here propagation is maintained due

to the delay in the production of v as shown in Fig. 2, compared to the production of v or the time at which v starts increasing in the standard model as shown in Fig. 3. Also the rate of recovery of the medium is made slower (in Fig. 2) compared to rate of recovery in the standard model where u goes to zero more rapidly (in Fig. 3).




Figure 1. Phase diagram illustrating the dynamics of the PDE-system with u and v recorded at the point $(40, 40)$ in a grid size of 220. Parameters: $a = 0.75, b = 0.06, \epsilon^{-1} = 13.5$. Shown are the nullclines: $v = u^3, u = u_{th}, u = 0$, and $u = 1$.




Figure 2. The time variation of u and v corresponding to one excursion along the phase diagram in Fig. 1.

3 Core Expansion

A spiral wave was initiated using the coarse gradient cross-field method, that is by setting $u = 0$ in the left half of the medium and $u = 1$ in the right half; $v = a/2$ in the upper half and $v = 0$ in the lower half. The subsequent evolution of the system at different times is shown in snapshots in Fig. 4. The simple rigid rotation of the spiral is destabilized because of the relatively high value of ϵ . The spiral tip defined as the intersection of the two isolines $u = 0.5$ and $v = 0.5u - b$, starts meandering, the spiral core expands as shown in Fig. 4, and the tip follows an outward motion along a spiraling




Figure 3. The time variation of u and v in the standard model.

path as shown in Fig. 5(left). The type of meandering is different for a different value of ϵ in Fig. 5(right). The meandering shown in Fig. 5(left) offers a demonstration that agrees with the prediction of the theory of non-local effects by Meron in the case of oscillatory recovery in excitable media. The type of recovery actually depends on the value of the control parameter ϵ . For low values of ϵ , perturbations near the spiral core are quenched by repulsive wave-fronts interactions in the monotonically recovering medium. If the value of ϵ is increased and for appropriate values for the other parameters, the tip undergoes this interesting outward motion along a path that looks like a spiral while the core grows in size due to oscillatory recovery to the resting state. The distance between the tip segment and the one ahead of it is determined by one of the maxima of the oscillatory tail. This corresponds to one of the minima of the excitation threshold.




Figure 4. Snapshots showing the core expansion with $dt = 0.052, dx = 0.51, L = 48$, grid size: 95. (Time intervals between snapshots are not equal).

In Fig. 5(left), the distance between the points denoting the tip positions increases as the tip moves outward implying that the tip motion is accelerated. In Fig. 4, the spiral core expands until spiral breakup occurs. This happens because the spiral period changes as the spiral drifts and meanders, until at some point within the excitable medium, it reaches the minimum period needed for plane wave propagation. This change in the spiral period as the tip moves outward and for-




Figure 5. Trajectory of the spiral tip defined as the intersection of the isolines $u = v = 0.5$, and following an outward spiral trajectory in (left) and meandering in (right). Parameters are the same as in Fig. 1 and Fig. 2 $\epsilon^{-1} = 18.0$.

ward is due to Doppler shift since the core is seen as the source of waves. This means that conduction would be blocked since the spiral rotates more rapidly than plane waves can propagate when the spiral period and the minimum period compatible with plane wave propagation merge for this critical value of ϵ [22]. The spiral becomes unstable and breaks into newly born broken waves which will soon evolve into spirals waves since they have broken ends. This leads to spatiotemporal chaos or irregularity within the excitable medium as shown in Fig. 6 where the time variations of the excitatory and the recovery variables are recorded at point (10, 10) in the excitable medium of size $L = 100$.




Figure 6. Time variation of the excitatory and the recovery variable recorded in the medium at the location (10, 10). The dotted one is the variation of v .

4 Traveling Waves

This phenomenon can be attributed to an unstable focus by considering a traveling wave solution of (1), $u(z) = u(x + ct)$ where c is the wave speed. Substituting this solution into (1) reduces the reaction-diffusion equations to the following ODE system:

$$\begin{aligned}
 dw/dt &= cw - (1/\epsilon)u(1-u)[u - ((b+v)/a)] \\
 du/dt &= w \\
 dv/dt &= (1/c)(u^3 - v),
 \end{aligned}
 \tag{2}$$

Using parameter values $a = 0.75$ and $b = 0.06$, $\varepsilon^{-1} = 13.5$, the numerical solution shown in Fig. 7 approaches the resting state in an oscillatory manner. For those values of the parameter, the system can have complex eigenvalues implying that the fixed point $(0, 0, 0)$ is an unstable focus for $c^2 - 4.32 < 0$. This condition is satisfied for some range of the spiral period as it can be seen in the dispersion curve in Fig. 8. That justifies the oscillatory behavior of the numerical solution of (2). For other values of the parameter ε , the solution approaches the resting state $(0, 0, 0)$ in a non-oscillatory manner if $\varepsilon^{-1} = 50.0$ for which the spiral rotates rigidly around a circular core. If $\varepsilon^{-1} = 20.0$, the recovery is also monotonic, but that does not necessarily imply rigid rotation. Actually, the spiral tip meanders following an epicycle-like orbit as shown in Fig. 5(right). On the other hand, if $\varepsilon^{-1} = 13.5$ or $\varepsilon^{-1} = 14.5$, the solution returns to the resting state in an oscillatory manner as seen in Fig. 7; the system undergoes a succession of super and subnormal periods until complete recovery is achieved. However, for $\varepsilon^{-1} = 14.5$, unlike the case for $\varepsilon^{-1} = 13.5$ (shown in Fig. 5 and Fig. 4) and despite the oscillatory type of recovery, core expansion does not occur and the tip does not follow an outward spiraling trajectory. It traces loops like those of an epicycle as the spiral wave rotates and drifts away. Actually, we found that the tip moves along a spiraling path in the range $13.0 < \varepsilon^{-1} < 13.8$. For $\varepsilon^{-1} > 13.9$, it meanders but not along a spiraling trajectory. Thus, we note the important conclusion that oscillatory recovery does not necessarily lead to core expansion and spiraling tip.




Figure 7. Recovering solitary traveling solution $u(z) = u(x + ct)$ of Eqs. (2), illustrating damped oscillatory ($\varepsilon^{-1} = 13.5, 14.5$) and monotonic recovery ($\varepsilon^{-1} = 20.0, 50.0$).

5 Dispersion Curve

This oscillatory recovery can be further investigated by writing the solution of the reaction-diffusion system as a superposition of two solitary waves with a small perturbation term R which vanishes in the limit of infi-

nite spacing between the two waves:

$$u(z) = u(z - z_1) + u(z - z_2) + R \quad (3)$$

where $z_1 = x_1 - ct$ and $z_2 = x_2 - ct$, x_1 and x_2 denote the waves positions. For large $|z_1|$ and $|z_2|$, the tail of the wave determines the manner in which the medium recovers to the resting state is: When the recovery is damped oscillatory, the tail of the wave $u(z) \propto e^{\eta z} \cos(\nu z + \psi)$; when it is monotonic, $u(z) \propto e^{\eta z}$. In both cases, the leading edge of the wave is assumed to be of the form $u(z) \propto e^{-\mu z}$. Equations for x_1 and x_2 are derived using the solvability conditions which remove singularities from R [Elphick *et al*, 1990]:

$$\frac{\delta x_1}{\delta t} = c + a_R e^{-\mu(x_1 - x_2)} \quad (4)$$

$$\frac{\delta x_2}{\delta t} = c + a_L e^{-\eta(x_1 - x_2)} \cos(\nu(x_1 - x_2) + \psi) \quad (5)$$

where c is the propagation speed of a solitary wave. The second term on the right hand side of (4) represents the effect of the second wave on the propagation of the first one. It is usually negligible in excitable media. The second term on the right hand side of (5) represents the effect exerted on the second wave by the refractory wake of the first one. Using (4) and (5), the spacing between the two waves $\lambda = x_1 - x_2$ obeys the equation:

$$\frac{d\lambda}{dt} = a_L e^{-\eta\lambda} \cos(\nu\lambda + \psi) \quad (6)$$

If $\nu \neq 0$, the excitable medium recovers in an oscillatory way. Then, according to (6) an infinite number of steady state solutions exist. This means that the distance between the wave-fronts takes one of possible fixed values.

Those oscillations in the way of recovery to the resting state would imply oscillations in the dispersion. This is shown here by considering the times when wave-fronts pass through a given location x . The solution of (5) is then approximated by widely spaced impulses:

$$u(x, t_i) = \sum_k u_k(t_i(x)) + R, \quad (7)$$

where $t_i(x)$ is the instant at which the i -th impulse is at x , and R is a small perturbation term which vanishes in the limit of infinite spacing between the waves. Using (7) in (1), we get

$$\frac{dt_i}{dx} = \frac{1}{c_0} + d' e^{\eta(t_i - t_{i-1})} \cos[\nu(t_i - t_{i-1}) + \psi] + b' e^{-\mu(t_{i+1} - t_i)} \quad (8)$$

where η is the rate at which the wave-fronts tail off. The second term on the right hand side of (8) represents the effect exerted on the i -th impulse by the refractory wake of the preceding impulse. The last term represents the effect of the succeeding impulse and is negligible in excitable media. The coefficients a' and b' require the evaluation of certain integrals which are not shown here. Let $t_i(x) = (x/c) + (i-1)T$, where T is the period of a constant speed wave-train. Then we get to leading order,

$$c = c_0 - c_0^2 a' e^{-\eta T} \cos[vT + \psi] \quad (9)$$

For $v \neq 0$ in (9), damped oscillations occur in the dispersion curve. For monotonic recovery ($v = 0, a > 0$), the wave speed is a monotonic increasing function of wave spacing. If $v = 0$ and $a < 0$, the recovery is said to be non-monotonic and could exhibit one supernormal period [Meron, 1991a], [Barkley, 1991]. In Fig. 8, there are damped oscillations in dispersion curve of the system (1) for $\epsilon^{-1} = 13.5$ and $\epsilon^{-1} = 14.5$. The first supernormal period during which the excitability is higher than that of the rest state is very pronounced. However, for $\epsilon^{-1} = 50.0$, as expected, the monotonically recovering system is characterized by a dispersion curve with monotonic increase in the propagation velocity until the limit set by the solitary wave velocity is reached.




Figure 8. Dispersion curve: Instantaneous wave speed as a function of period for $\epsilon^{-1} = 13.5, 14.5, 50.0$ respectively from bottom to top, computed by repetitively stimulating at one of the ends of an open line.

6 Resonant Drift

Furthermore, we propose the Feedback Resonant Drift Method to control the spatiotemporal chaotic behavior or irregularity shown in Fig. 6. A recording point is set at the upper right corner of the medium. Each time the recorded variable u crosses a certain threshold value 0.8, a small amplitude non-localized stimulation is applied with amplitude 0.26 t.u., by adding it to the

right hand side of (1). We verified that the amplitudes needed to eliminate all wave activity were between 2% and 11% of the amplitude of a single shock that would eliminate all wave activity. This single shock (with defibrillation amplitude 0.055) would represent the single shock defibrillation threshold. In Fig. 9, the system is left with only one spiral to be eliminated at the unexcitable boundaries before breakup occurs again and generates further spiral waves. The directed drift due to the perturbation pulses did not lead to elimination at the upper boundary but at the left side of the medium. In Fig. 10, the tip path is shown as the repetitive perturbations are applied until the path ends at the left boundary.




Figure 9. Snapshots illustrating the evolution of the system under perturbation pulses with amplitude 0.0057 and pulse duration 0.26 t.u.




Figure 10. Tip trajectory when perturbations are applied.

This oscillatory behavior and the occurrence of supernormal periods in the dispersion curve were observed in wave train solutions of the one-dimensional FitzHugh-Nagumo model [Rinzel and Maginu, 1984]. But, no core expansion has ever been reported before. Also, the observation of expanding cores and spiraling tips here answers the query of Meron [Meron, 1991b], [Meron, 1989], [Meron, 1991a] and Winfree [Winfree, 1991], [Winfree, 1990] about the possible observation of core expansion and oscillations in the dispersion curve. Using FitzHugh-Nagumo kinetics with parameters chosen such that the equilibrium point is nearly

a center, Winfree showed that the medium can then support two stable rotors of different periods. The dispersion curve exhibited a damped oscillatory behavior. However, core expansion was not observed, and meandering along a spiraling path was not obtained.

7 Conclusion

This occurrence of 'supernormal' periods of excitability during which the threshold of excitation is diminished was reported in electrophysiological measurements in stimulated cardiac muscle [Spear and Moore, 1974]. The current that was needed to re-excite the Purkinje fibers was reduced. We could attribute it to the faster recovery of the threshold potential compared to the slower recovery of the action potential that we have seen here. We have also verified that than a smaller additional depolarization is needed to reach the threshold potential and it was brought about by a weaker depolarizing current.

Our results would imply that core expansion could be one possible route to spiral breakup. We also offered one effective way to eliminate wave activity, by applying feedback perturbations. This represents one way that would help to eliminate cardiac arrhythmias or fibrillation by causing spiral waves to move outward toward the unexcitable boundaries or non-cardiac tissue.

8 Acknowledgments

This project has been supported by a grant from the CNRS.

The author would like to thank Prof. A.V. Holden.

References

- Barkley, D. (1991) A model for fast computer simulation of waves in excitable media. *Physica D*, **49**, pp. 61–70.
- Barkley, D., Kness, M., and Tuckerman, L.S. (1990) Spiral-wave dynamics in a simple model of excitable: The transition from simple to compound rotation. *Phys. Rev. A*, **42**, pp. 2489–2492.
- Special issue on dynamics in cardiac tissue. (2002) *Chaos*, **12** (3).
- Courtemanche, M., Glass, L., and Kenner, J.P. (1993) Instabilities of a propagating pulse in a ring of excitable media. *Phys. Rev. Lett.*, **70**, pp. 2182–2185.
- Elphick, C., Meron, E., Rinzel, J., and Spiegel, E.A. (1990) Impulse patterning and relaxational propagation in excitable media. *J. Theor. Biol.*, **146**, pp. 249–268.
- Imbihl, R., and Ertl, G. (1995) Oscillatory kinetics in heterogeneous catalysis. *Chem. Rev.*, **95**, pp. 697–733.
- Kapral, R., and Showalter, K. (Eds.) (1994) *Chemical Waves and Patterns*. Dordrecht: Kluwer.
- Karma, A. (1994) Electrical alternans and spiral wave breakup in cardiac tissue. *Chaos*, **4**, pp. 461–472.
- Markus, M., and Stavridis, K. (1994) Observation of chemical turbulence in the Belousov-Zhabotinsky reaction. *Int. J. Bifurc. Chaos*, **4**, pp. 1233–1243.
- Meron, E. (1989) Nonlocal effects in spiral waves. *Phys. Rev. Lett.*, pp. 684–687.
- Meron, E. (1991a) The role of curvature and wavefront interactions in spiral-waves interactions. *Physica D*, **49**, pp. 98–106.
- Meron, E. (1991b) The Effect of wavefront interactions on pattern formation in excitable media. *Nonlinear Wave Processes in Excitable Media*. A.V. Holden et al. (Eds.) Plenum, pp. 145–153.
- Murray, J. (1989) *Mathematical Biology*. Berlin: Springer, 1989.
- Nagy-Ungvarai, Zs., and Muller, S.C. (1994) Characterization of wave front instabilities in the Belousov-Zhabotinsky reaction: an overview. *Int. J. Bifurc. Chaos*, **4**, pp. 1257–1264.
- Panfilov, A.V., and Hogeweg, P. (1995) Turbulence in three-dimensional excitable media. *Science*, **270**, pp. 1223–1224.
- Rinzel, J., and Maginu, K. (1984) Kinematic analysis of wave pattern formation in excitable media. *Nonequilibrium Dynamics in Chemical Systems*. Vidal, C., and Pacault, A. (Eds.) Springer-Verlag, pp. 107–113.
- Sabbagh, H. (2000) Stochastic properties of autowave turbulence elimination. *Chaos Solitons & Fractals*, **11**, pp. 2141–2148.
- Sabbagh, H. (2009) Spatiotemporal chaos and break up of spiral waves due to core expansion in excitable media. *Journal of Applied Functional Analysis*, **4**(3), pp. 623–635.
- Spear, J.F., and Moore, E.N. (1974) Supernormal excitability and conduction in the His-Purkinje system of the dog. *Circ. Res.*, **35**, pp. 782–792.
- Tyson, J.J., and Keener, J.P. (1988) Singular perturbation theory of traveling waves in excitable media (a review). *Physica D*, **32**, pp. 327–361.
- Winfree, A.T. (1987) *When Time Breaks Down*. Princeton Univ. Press: New Jersey.
- Winfree, A.T. (1990) Discrete Spectrum of rotor periods in an excitable medium. *Phys. Lett. A*, **149**, pp. 203–206.
- Winfree, A.T. (1991) Alternative stable rotor in an excitable medium. *Physica D*, **49**, pp. 125–140.
- Winfree, A.T. (1994) Electrical turbulence in three dimensional heart muscle. *Science*, **266**, pp. 1003–1006.
- Zykov, V.S. (1986) Cycloid circulation of spiral waves in an excitable media. *Biophysics*, **31**(5), pp. 940–944.
- Zykov, V.S. (1987) Kinematics of the non-steady circulation of helical waves in excitable media. *Biophysics*, **32**(2), pp. 365–369.
- Zykov, V.S., Steinbock, O., and Muller, S.C. (1994) External forcing of spiral waves. *Chaos*, **4**(3), pp. 509–518.