

## THE DYNAMICS OF EEG BRAIN PATTERNS IN THE PROCESS OF BISTABLE IMAGE PERCEPTION

**Anastasiya Runnova**

Department of  
Automation, Control, Mechatronics,  
Saratov State Technical University,  
Russia  
anefila@gmail.com

**Vadim Grubov**

Department of  
Automation, Control, Mechatronics,  
Saratov State Technical University,  
Russia  
vvgrubov@gmail.com

**Alexey Koronovskii**

Faculty of Nonlinear Processes,  
Saratov State University,  
Russia  
alexey.koronovskii@gmail.com

**Maksim Zhuravlev**

Faculty of Nonlinear Processes,  
Saratov State University,  
Russia  
zhuravlevmo@gmail.com

**Alexander Hramov**

Department of  
Automation, Control, Mechatronics,  
Saratov State Technical University,  
Russia  
hramovae@gmail.com

**Alexander Pisarchik**

Center for Biomedical Technology  
Technical University of Madrid  
Spain  
alexander.pisarchik@ctb.upm.es

### Abstract

In the paper we study the dynamics of the complex patterns in human EEG data during a psychophysiological experiment by stimulating cognitive activity with the perception of ambiguous object. A new method based on the calculation of the maximum energy component for the continuous wavelet transform (skeletons) is proposed. The paper presents the processing results of experimental data for 20 male volunteers. Skeleton analysis allows us to identify specific patterns in the EEG data set, appearing in the perception of ambiguous objects. Thus, it becomes possible to diagnose some cognitive processes associated with the concentration of attention and recognition of complex visual objects. In addition, with enough confidence we define the periods of relaxation, appearing in the intervals between repeated moments of perception of ambiguous objects. Studies have shown that for a sufficiently long discontinuous perception of an object process appears at a different rate for various volunteers, but with a similar dynamics of learning to "rest" (relaxation) between perceptions of ambiguous images.

### Key words

Multistability, bistable image, perception, EEG, continuous wavelet transform

### 1 INTRODUCTION

Nowadays, the study of brain dynamics in cognitive activity attracted much attention of researchers. Such studies often used electroencephalography, because this method is non-invasive and does not require significant limitations volunteer mobility, nor for costs. This article is devoted to the study of the popular des-

tinuations in cognitive perception of ambiguous images. The investigations of nonlinear processes in the brain neural network during perception of ambiguous (the so-called *bi-* and *multistable*) images are very important for the understanding of both the visual recognition of objects and the decision-making process.

Images of this type have been the object of research for psychologists for a long time [Leopold, Logothetis, 1999; Sterzer, Kleinschmidt, Rees, 2009]. Recently, ambiguous images awoke interest of physicists and mathematicians [Pisarchik, Jaimes-Reategui, Magallon-Garcia et al., 2014; Runnova, Hramov, Grubov et al., 2016]. Despite of considerable efforts of many researchers, the main mechanisms underlying interpretation of a multistable image are not well understood. At present, perception is known to be a result of nonlinear processes which take place in the distributed neural network of occipital, parietal and frontal regions of the brain cortex [Sterzer, Kleinschmidt, Rees, 2009; Tong, Meng, Blake, 2006].

The perception of ambiguous (bistable) images was thoroughly investigated in the last decade. The most popular bistable images are Rubin vase, Mach bands, Rorschach test, Boring's old/young woman illusion, Necker cube, etc. [Huguet, Rinzel, Hup, 2014; Gigante, Mattia, Braun et al., 2009; Moreno-Bote, Rinzel, Rubin, 2007; Merk, Schnakenberg, 2002; Aks, Sprott, 2003; Ratcliff, Smith, 2004; Heekeren, Marrett, Ungerleider, 2008]. From a mathematical point of view, visual perception of bistable images comprises two metastable states, each with a duration that varies from seconds to tens of seconds. One of the perceived images usually dominates over the other for a relatively long period of time [Wang, 2012; Pearson, Raskevičius, Bays et al., 2014].

These fundamental results are important to create an objective paradigm for data processing in cognitive research, because the application of traditional neurophysiological methods solely, such as, expert estimations and simple measures of the signal amplitude, may simplify our understanding of the process. Therefore, the fine and short-lasting process might be overlooked. The development of standardized methods has an important practical application in neurophysiological experiments, including data acquisition, evaluation, analysis and processing of results in already recorded data, as well as in real time, and it is still an open field for research activity. International research in neurophysiological studies readily engages in the modeling of cognitive and pathological processes in the brain neuronal network using nonlinear dynamics tools [Rabinovich, Varona, Selverston et al., 2006; Hramov, Koronovskii, Midzyanovskaya et al., 2006; Sitnikova, Hramov, Grubov et al., 2012; Hramov, Koronovskii, Makarov et al., 2015; van Luijtelea, Luttjohann, Makarov et al., 2016; Hramov, Sitnikova, Pavlov et al., 2015].

## 2 STUDY METHODS

### 2.1 Experiment

The experimental studies were performed in accordance with the ethical standards of the World Medical Association [World medical association, 2000]. Twenty healthy subjects from a group of unpaid male volunteers, between the ages of 19 and 30 with a normal visual acuity participated in the experiments. The purpose of this experiment is the study of multi-channel EEG data registration in the unconscious decision on ambiguous image interpretation.

In our physiological experiment with EEG activity registration we used a set of images based on a well-known bistable object, the Necker cube [Necker, 1832], as a visual stimulus. This is a cube with transparent faces and visible ribs; an observer without any perception abnormalities treats the Necker cube as a 3D-object thanks to the specific position of the cube ribs. Bistability in perception consists in the interpretation of this 3D-object as to be oriented in two different ways, in particular, if the different ribs of the Necker cube are drawn with different intensity. In our experimental works we have used the Necker cube images with varying parameter  $I$  to be the brightness of the cube wires converging in the right upper inner corner (Fig. 1). The brightness of the wires converging in the left lower inner corner is defined as  $(1 - I)$ .

We demonstrated the Necker cube images with different wireframe contrasts for a short time, each lasting between 1.5 and 2.0 seconds, interrupted by a background abstract picture for 5.0 – 5.5 seconds. The subject was instructed to press the left or right key depending on his/her interpretation of the projection being observed at each demonstration. The use of the background abstract images allows the neutraliza-

tion of possible negative secondary effect of the previous Necker cube image. The whole experiment lasted about 40 min for each patient.




Figure 1. Examples of distinct Necker cube images with different wireframe contrasts characterized by control parameter  $I$  and fragments of EEG data. These curves are plotted according to the data in occipital area channels volunteer number 1.

During the experiment we exhibited the pictures of the Necker cube randomly, each for about 100 times, and recorded brain activity with multi-channel EEG. As a tool for EEG recording we used the electroencephalograph–recorder Encephalan-EEGR-19/26 (Russia) with multiple EEG channels and the two-button input device. To study EEGs the monopolar registration method and the classical ten-twenty electrode system were used. Figure 1 shows an example of a typical EEG data set from EEG registration channels of occipital area. It seems occipital region associated with cognitive processes of perception of complex spatial objects, which include the Necker cube.

### 2.2 Wavelet-based method

In our work we used continuous wavelet transform (CWT) [Hramov, Koronovskii, Makarov et al., 2015; Koronovskii, Ponomarenko, Prokhorov et al., 2007; Hramov, Koronovskii, Ponomarenko et al., 2006; Hramov, Koronovskii, Ponomarenko et al., 2007; Pavlov, Hramov, Koronovskii et al., 2012] for time-frequency analysis of oscillatory patterns in EEG. CWT is a convolution of investigated signal  $x(t)$  (EEG signal in our case) and a set of basic functions  $\varphi_{s,\tau}$ :

$$W(x, \tau) = \int_{-\infty}^{\infty} x(t) \varphi_{s,\tau}^* dt \quad (1)$$

Each basic function from this set can be obtained from one function  $\varphi_0$ , the so-called mother wavelet, by fol-

lowing transform:

$$\varphi(s, \tau) = \frac{1}{\sqrt{s}} \varphi_0 \left( \frac{t - \tau}{s} \right) \quad (2)$$

In equation (2)  $\varphi_0$  — mother wavelet,  $s$  — time scale, which determines extension or compression of initial mother function,  $\tau$  — time shift of wavelet transform.

There are a lot of different mother wavelets that find a use according to the problems of the current study. In present work we used CWT with Morlet mother wavelet with parameter  $\omega_0 = 2\pi$  [Ovchinnikov, Lutjtjohann, Hramov et al., 2010; Ovchinnikov, Hramov, Lutjtjohann et al., 2011]:

$$\varphi_0(\eta) = \pi^{-\frac{1}{4}} e^{j\omega_0\eta} e^{-\frac{\eta^2}{2}} \quad (3)$$

According to papers [Hramov, Koronovskii, 2004; Hramov, Koronovskii, Ponomarenko et al., 2006; Sitnikova, Hramov, Koronovskii et al., 2009; Luijtelaar, Hramov, Sitnikova et al., 2011; Sitnikova, Hramov, Grubov et al., 2014] the Morlet wavelet is one of the most effective in analysis of complex experimental signals of biological nature (including EEG data) because of its optimal time-frequency resolution.

In present work intrinsic frequency dynamics was investigated using “skeletons” of wavelet surfaces [Sitnikova, Hramov, Grubov et al., 2014; Sitnikova, Grubov, Hramov et al., 2011; Hramov, Kharchenko, Makarov et al., 2016]. The “skeletons” of wavelet surfaces are constructed to extract dominant EEG frequencies and determine the evolution of oscillatory patterns in EEG data. In this paper, we focused on the processing of EEG data recorded in the occipital region (see Fig. 1). First, the momentary wavelet energy distribution  $E_i(f_s, t_0)$  was constructed for some time moment  $t_0$ .

$$E_i(f_s, t_0) = |W(f_s, t_0)|^2 \quad (4)$$

Then the function  $E_i(f_s, t_0)$  was examined for the presence of local maximum  $E_{max}$ . If several local maxima  $E_{max,k}$  were detected in  $E_i(f_s, t_0)$ , then the highest maximum was selected and its frequency was considered as dominant frequency of oscillatory pattern at given time moment  $t_0$ . In order to construct full “skeleton” of wavelet surface the procedure described above was repeated consequently for all points in time series of given EEG signal.

### 3 RESULTS

In our experimental studies we recorded EEG data during the sufficiently regular ambiguous perception of complex objects. Between the two moments of the perception of ambiguous object associated with a spatial imagination Necker cube, a pause is enough to

restore the background activity is not associated with the cognitive process of spatial modeling. At the moment, we are not interested in causes, for example, the phenomenon of cognitive noise, forcing Necker cube with intensity  $I$  perceived as the “left” and “right” [Pisarchik, Bashkirtseva, Ryashko, 2015]. However, we directly focused on the processes that occur in the perception and the internal filling volume of a two-dimensional object Necker ambiguous image. The visual analysis of skeleton dynamics is not enough for math description of the brain neurophysiological processes, and we propose a method for an estimation of level and nature of beta-activity for occipital region in the human brain. For this we consider the main characteristic of the frequency range of 25–35 Hz, well traced throughout the duration of the processes associated with the recognition of ambiguous images for the majority of volunteers. Note that one volunteer with displacement of the pattern frequency to 30–38 Hz was found in the processing data. However, it is clear that this fact is easily detected and can be corrected adaptation algorithm.

Now we are limiting our processing of multi-channel EEG data exceptionally occipital area of a brain electrical activity registration, in particular O1, O2, P3, P4, Pz, C3, C4 and Cz channels of the classical ten-twenty electrode system. We introduce a numerical criterion for the presence of alpha-rhythm in every time moment: the values of first two skeletons are in the range of 30 to 40 Hz in this time. In other words, if the condition is satisfied for a particular channel of the occipital region, the alpha-criterion  $\beta$  is equivalent to the constant, and otherwise  $\beta$  is zero, then  $\beta$  – criterion:

$$\beta_i = \begin{cases} 1 & \text{if } 30 < f_{1,2}^i < 40 \\ 0 & \text{in other cases} \end{cases}$$

In Figure 2 a shows the result of calculation of the proposed specifications at any given time for the channel Pz. This EEG-signal registration is corresponding to the electricity activity of the interhemispheric slot in human brain. However, consideration of the signal for each EEG channel is not convenient and leads to the accumulation of errors. We have tried to average characteristics of excitable beta-activity for the whole occipital region. Note that in this case we consider the spatial and temporal dynamics of occurring beta-pattern correlated with the cognitive process of recognition of a complex ambiguous object. In this case the presence of the beta-rhythm criterion for each channel, it can simple sum the “0” and “1”:

$$\beta_{8i} = \sum \begin{cases} 1 & \text{if } 30 < f_{1,2}^{O1} < 40 \text{ or } 0 \text{ in other cases} \\ 1 & \text{if } 30 < f_{1,2}^{O2} < 40 \text{ or } 0 \quad \dots \\ 1 & \text{if } 30 < f_{1,2}^{P3} < 40 \text{ or } 0 \quad \dots \\ 1 & \text{if } 30 < f_{1,2}^{Pz} < 40 \text{ or } 0 \quad \dots \\ 1 & \text{if } 30 < f_{1,2}^{P4} < 40 \text{ or } 0 \quad \dots \\ 1 & \text{if } 30 < f_{1,2}^{C3} < 40 \text{ or } 0 \quad \dots \\ 1 & \text{if } 30 < f_{1,2}^{Cz} < 40 \text{ or } 0 \quad \dots \\ 1 & \text{if } 30 < f_{1,2}^{C4} < 40 \text{ or } 0 \quad \dots \end{cases}$$

In Figure 2 b we show the calculation result of the characteristics of  $\beta$ -activity for the whole of the occipital region of the brain. However, we paid attention to the existence of certain regularities in the distribution of  $\beta$ -pattern in the occipital region, directly related to the nature of cognitive activity.




Figure 2. (a) The dependence of the beta-criterion  $\beta$  over time for Pz channel. (b) The dependence of the beta-criterion  $\beta_{8i}$  for brain occipital region.

#### 4 CONCLUSION

In this paper we have considered the technique for study evolution of the complex patterns in human EEG data during a psychophysiological experiment by stimulating cognitive activity with the perception of ambiguous object. The new method based on continuous wavelet transform allows to estimate the energy contribution of various components in the general and partial dynamics of the electrical activity for the projections of various areas of the brain.

The results of these studies appear promising for further study of the dynamics and the activity of the cerebral cortex in cognitive processes of various kinds. The technique is based on the calculation of the wavelet skeleton, it is universal for the study of various processes. Furthermore, this approach is highly customizable to individual features volunteers that allows the theoretical possibility of that using in the biofeedback systems.

#### 5 ACKNOWLEDGMENTS

This work has been supported by the Russian Science Foundation (Grant No. 16-12-10100).

#### References

- Leopold, D. A. and Logothetis, N. K. (1999) Multistable phenomena: changing views in perception. *Trends in Cognitive Sciences*, **3**(7), pp. 254–264.
- Sterzer, P., Kleinschmidt, A., and Rees, G. (2009) The neural bases of multistable perception. *Trends in Cognitive Sciences*, **13**(7), pp. 310–318.
- Pisarchik, A. N., Jaimes-Reategui, R., Magallón-García, C. D. A., and Castillo-Morales, C. O. (2014) Critical slowing down and noise-induced intermittency in bistable perception: bifurcation analysis. *Biological Cybernetics*, **108**(4), pp. 397–404.
- Runnova, A. E., Hramov, A. E., Grubov, V. V., Koronovskii, A. A., Kurovskaya, M. K., and N., P. A. (2016) Theoretical background and experimental measurements of human brain noise intensity in perception of ambiguous images. *Chaos, Solitons and Fractals*, **93**, pp. 201–206.
- Tong, F., Meng, M., and Blake, R. (2006) Neural bases of binocular rivalry. *Trends in Cognitive Sciences*, **10**(11), pp. 502–511.
- Huguet, G., Rinzel, J., and Hupé, J.-M. (2014) Noise and adaptation in multistable perception: Noise drives when to switch, adaptation determines percept choice. *Journal of Vision*, **14**(3), pp. 1–24.
- Gigante, G., Mattia, M., Braun, J., and Giudice, P. D. (2009) Bistable perception modeled as competing stochastic integrations at two levels. *PLoS Computational Biology*, **5**(7), e1000430.
- Moreno-Bote, R., Rinzel, J., and Rubin, N. (2007) Noise-induced alternations in an attractor network model of perceptual bistability. *Journal of Neurophysiology*, **98**, pp. 1125–1139.
- Merk, I. and Schnakenberg, J. (2002) A stochastic model of multistable visual perception. *Biological Cybernetics*, **86**, pp. 111–116.
- Aks, D. J. and Sprott, J. C. (2003) The role of depth and  $1/f$  dynamics in perceiving reversible figures. *Nonlinear Dynamics, Psychology, and Life Sciences*, **7**(2), pp. 161–180.
- Ratcliff, R. and Smith, P. L. (2004) A comparison of sequential sampling models for two-choice reaction time. *Psychological Review*, **111**(2), pp. 333–367.
- Heekeren, H. R., Marrett, S., and Ungerleider, L. G. (2008) The neural systems that mediate human perceptual decision making. *Nature Reviews Neuroscience*, **9**, pp. 467–479.
- Wang, X.-J. (2012) Neural dynamics and circuit mechanisms of decision-making. *Current Opinion in Neurobiology*, **22**(6), pp. 1039–1046.
- Pearson, B., Raskevicius, J., Bays, P. M., Pertsov, Y., and Husain, M. (2014) Working memory retrieval as a decision process. *Journal of Vision*, **14**(2).
- Rabinovich, M. I., Varona, P., Selverston, A. I., and Abarbanel, H. D. I. (2006) Dynamical principles in neuroscience. *Rev. Mod. Phys.*, **78**, pp. 1213–1265.
- Hramov, A. E., Koronovskii, A. A., Midzyanovskaya, I. S., Sitnikova, E., and Rijn, C. M. (2006) On-off intermittency in time series of spontaneous parox-

- ysmal activity in rats with genetic absence epilepsy. *Chaos*, **16**, 043111.
- Sitnikova, E., Hramov, A. E., Grubov, V. V., Ovchinnikov, A. A., and Koronovsky, A. A. (2012) On-off intermittency of thalamo-cortical oscillations in the electroencephalogram of rats with genetic predisposition to absence epilepsy. *Brain Research*, **1436**, pp. 147–156.
- Hramov, A. E., Koronovskii, A. A., Makarov, V. A., Pavlov, A. N., and Sitnikova, E. (2015) [*Wavelets in Neuroscience*], Springer Series in Synergetics, Springer, Heidelberg, New York, Dordrecht, London.
- van Luijtelaar, G., Luttjohann, A., Makarov, V. V., Maksimenko, V. A., Koronovskii, A. A., and Hramov, A. E. (2016) Methods of automated absence seizure detection, interference by stimulation, and possibilities for prediction in genetic absence models. *Journal of Neuroscience Methods*, **260**, pp. 144–158.
- Hramov, A. E., Sitnikova, E. Yu., Pavlov, A. N., Grubov, V. V., Koronovskii, A. A., Khramova, M. V. (2015) Time-frequency dynamics during sleep spindles on the EEG in rodents with a genetic predisposition to absence epilepsy (WAG/Rij rats). *Proc. SPIE*, **9448**, 94481P.
- World medical association (2000) Declaration of helsinki: ethical principles for medical research involving human subjects., *The Journal of the American Medical Association*, **284**(23), pp. 3043–3045.
- Necker, L. A. (1832) Observations on some remarkable phenomena seen in Switzerland; and an optical phenomenon which occurs on viewing of a crystal or geometrical solid. *Philos. Mag.*, **3**, pp. 329–343.
- Koronovskii, A. A., Ponomarenko, V. I., Prokhorov, M. D., and Hramov, A. E. (2007) Method of studying the synchronization of self-sustained oscillations using continuous wavelet analysis of univariant data. *Technical Physics*, **52**(9), pp. 1106–1116.
- Hramov, A. E., Koronovskii, A. A., Ponomarenko, V. I., and Prokhorov, M. D. (2006) Detecting synchronization of self-sustained oscillators by external driving with varying frequency. *Phys. Rev. E*, **73**, 026208.
- Hramov, A. E., Koronovskii, A. A., Ponomarenko, V. I., and Prokhorov, M. D. (2007) Detection of synchronization from univariate data using wavelet transform. *Phys. Rev. E*, **75**(5), 056207.
- Pavlov, A. N., Hramov, A. E., Koronovskii, A. A., Sitnikova, Y. E., Makarov, V. A., and Ovchinnikov, A. A. (2012) Wavelet analysis in neurodynamics. *Physics-Uspeski*, **55**(9), pp. 845–875.
- Ovchinnikov, A. A., Luttjohann, A., Hramov, A. E., and Luijtelaar van, G. (2010) An algorithm for real-time detection of spike-wave discharges in rodents. *Journal of Neuroscience Methods*, **194**, pp. 172–178.
- Ovchinnikov, A. A., Hramov, A. E., Luttjohann, A., Koronovskii, A. A., and van Luijtelaar, E. L., (2011) Method for diagnostics of characteristic patterns of observable time series and its real-time experimental implementation for neurophysiological signals. *Technical Physics*, **56**(1), pp. 1–7.
- Hramov, A. E. and Koronovskii, A. A. (2004) An approach to chaotic synchronization. *Chaos*, **14**(3), pp. 603–610.
- Sitnikova, E., Hramov, A. E., Koronovskii, A. A., and Luijtelaar, E. L., (2009) Sleep spindles and spike-wave discharges in EEG: Their generic features, similarities and distinctions disclosed with fourier transform and continuous wavelet analysis. *Journal of Neuroscience Methods*, **180**, pp. 304–316.
- van Luijtelaar, E. L. M., Hramov, A. E., Sitnikova, E., and Koronovskii, A. A., (2011) Spike-wave discharges in WAG/Rij rats are preceded by delta and theta precursor activity in cortex and thalamus. *Clinical Neurophysiology*, **122**, pp. 687–695.
- Sitnikova, E., Hramov, A. E., Grubov, V., and Koronovsky, A. A., (2014) Time-frequency characteristics and dynamics of sleep spindles in WAG/Rij rats with absence epilepsy. *Brain Research*, **1543**, pp. 290–299.
- Sitnikova, E. Yu., Grubov, V. V., Hramov, A. E., and Koronovskii, A. A., (2011) Structure of EEG sleep spindles in rats with genetic predisposition to absence epilepsy (WAG/Rij). *Journal of higher nervous activity*, **62**(6), pp. 733–744.
- Hramov, A. E., Kharchenko, A. A., Makarov, V. V., Khramova, M. V., Koronovskii, A. A., Pavlov, A. N., and Dana, S. K. (2016) Analysis of the characteristics of the synchronous clusters in the adaptive Kuramoto network and neural network of the epileptic brain. *Proc. SPIE*, **9917**, 991725.
- Pisarchik, A. N., Bashkirtseva, I. A., and Ryashko, L. B. (2015) Controlling bistability in a stochastic perception model. *Eur. Phys. J. Special Topics*, **224** (8), pp. 1477–1484.